

Cross-linking of micelles by gemini surfactants

Prabal K. Maiti^{1,*} and Kurt Kremer

Max Planck Institute for Polymer Research,

Postfach 3148, 55021 Mainz, Germany

Oliver Flimm, Debashish Chowdhury[†] and Dietrich Stauffer

Institute for Theoretical Physics,

University of Cologne, 50923 Köln, Germany

We investigate the effects of gemini surfactants, telechelic chain and lipids on the nature of micelles formed by conventional single-tail surfactants in water by carrying out Monte Carlo simulations. In a mixture of gemini and single-tail surfactants in water we find direct evidence of micelles of predominantly single-tail surfactants some of which are dynamically cross-linked by gemini surfactants when the concentrations of the geminis is only a few mole percent and their spacers are *hydrophilic*. In contrast, mixtures of lipids and single-tail surfactants in water form only isolated micelles, each consisting of a mixture of both species, without cross-links.

Running title: Mixed and cross-linked micelles

PACS Numbers: 68.10.-m, 82.70.-y

I. INTRODUCTION:

Amphiphilic molecules are not only important constituents of bio-materials, but also find wide-ranging industrial applications [1]. These surface-active agents are also referred to as surfactants. Many surfactants consist of a single "hydrophilic head" connected to a single "hydrophobic tail" whereas some other surfactants, e.g., phospholipids, are made of two hydrophobic tails both of which are connected to the same hydrophilic head (fig. 1). In contrast, *gemini* surfactants [2–8] consist of two single-tail surfactants whose heads are connected by a "spacer" chain (fig.1). The spacer in gemini surfactants may be hydrophilic or hydrophobic [9]. Because of their "water-loving" heads and "water-hating" tails, surfactant molecules form "*self-assemblies*" (i.e., supra-molecular aggregates), such as micelles, vesicles, etc., [10] in aqueous media. For example, micelles are formed when the concentration of the surfactants in water exceeds what is known as the "critical" micellar concentration (CMC). Some of the unusual properties of surfactants, which are exploited in their wide-ranging industrial applications, are crucially influenced by the nature of their aggregation in aqueous environments.

In this paper we address the following question: what are the effects of a few mole percent of gemini surfactants, mixed with single-tail surfactants, on the nature of micellar aggregates in water? By carrying out Monte Carlo (MC) simulations of a mixture of gemini and single-tail surfactants in water we show that in such a system, in equilibrium, micelles of predominantly single-tail surfactants are physically (i.e., non-covalently) "cross-linked" by geminis when the spacers are hydrophilic and the relative concentration of the geminis is only a few mole percent. The direct evidence emerging from the instantaneous snapshots of the system in our simulations may settle a controversy as to the interpretation of indirect experimental indications [11–13]. Moreover, we find lipids (i.e., double-tail surfactants) fail to crosslink micelles of single-tail surfactants; instead, isolated micelles, each consisting of a mixture of lipids and single-tail surfactants, form spontaneously in water. Furthermore, we also show that "telechelic chain" [14,15], another class of amphiphilic molecules, succeeds

less often than the geminis in cross-linking micelles.

II. THE MODEL

In the spirit of lattice gas models, we model the fluid under investigation as a simple cubic lattice of size $L_x \times L_y \times L_z$. A water molecule occupies a single lattice site. The surfactants are modelled as short chains on the lattice. The "primary structure" of a single-chain surfactant is defined by the symbol [16–18] $\mathcal{T}_m \mathcal{N}_p \mathcal{H}_q$ where $\mathcal{T}$ denotes *tail*, $\mathcal{H}$ denotes *head* and $\mathcal{N}$ denotes the 'liaison' or neutral part of the surfactants. The integers m, p and q denote the lengths of the corresponding pieces, giving a chain of length $\ell = (m + p + q)$. Bernardes [19] extended this to a microscopic lattice model of lipids in water. The "primary structure" of lattice model of lipids can be described by the symbol $\mathcal{T}_m \mathcal{N}_p \mathcal{H}_q \mathcal{N}_p \mathcal{T}_m$. In terms of the same symbols, the gemini surfactant can be represented [20] by the symbol $\mathcal{T}_m \mathcal{N}_p \mathcal{H}_q \mathcal{S}_n \mathcal{H}_q \mathcal{N}_p \mathcal{T}_m$ where n is the number of lattice sites constituting the spacer represented by the symbol $\mathcal{S}$. Finally, a microscopic lattice model of "telechelic" chains can be represented by the symbol $\mathcal{T}_1 \mathcal{N}_p \mathcal{H}_n \mathcal{N}_p \mathcal{T}_1$. In other words, a telechelic chain can be viewed as a gemini surfactant with hydrophilic spacer and the shortest possible hydrophobic tails.

Using these lattice models of surfactants in water, in this paper we investigate the effects of small mole fractions of geminis (or lipids or telechelic chains) on the micellar aggregates formed by the single-tail surfactants. We carry out Monte Carlo (MC) simulations of a mixture of single-tail and gemini surfactants in water for $p = q = 1$ and for tail length $m = 4$ to $m = 12$ and spacer length $n = 0$ to $n = 14$. We shall refer to each site on the surfactants as a *monomer*. Note that in this type of models [21] the "water-loving" head group is assumed to be "water-like".

The above mentioned model of single-chain surfactants in aqueous media [22] has been reformulated [16,17] in terms of Ising-like variables, in the same spirit in which a large number of simpler lattice models had been formulated earlier [23] for the convenience of calculations. In this formulation, a classical Ising spin-like variable S is assigned to each

lattice site; $S_i = 1$ (-1) if the i -th lattice site is occupied by a water (oil) molecule. If the j -th site is occupied by a monomer belonging to a surfactant then $S_j = 1, -1, 0$ depending on whether the monomer at the j th site belongs to head, tail or neutral part (see fig.2). The monomer-monomer interactions are taken into account in analogy with the ferromagnetic nearest-neighbour interaction in Ising magnets. Thus, the thermal Boltzmann probability for a configuration is proportional to $\exp(-H/k_B T)$ and the Hamiltonian for the system is given by the standard form

$$H = -J \sum_{\langle ij \rangle} S_i S_j. \quad (1)$$

The temperature T of the system is measured in the units of J/k_B where J denotes the strength of the interactions between the spin-like variables S on nearest-neighbour sites and k_B is the Boltzmann constant (while varying T we assume J to be constant). Moreover, in order to investigate the role of the chain stiffness we have used a chain bending energy [24]; every bend of a spacer, by a right angle at a lattice site, is assumed to cost an extra amount of energy $K(> 0)$.

This type of microscopic lattice models is very useful for investigating the generic qualitative features of a class of surfactants rather than for a quantitative prediction of any dynamic property of any specific material; for the latter purpose molecular dynamics simulations of molecular model of the specific surfactant is more appropriate, although much more computer time consuming.

In order to investigate the spontaneous formation and morphology of supra-molecular aggregates, we initially disperse the model gemini surfactants and single tail surfactants randomly in a $L_x \times L_y \times L_z$ system which contains only water other than the surfactants. Since we are interested in the effects of gemini surfactants on micellar aggregates formed by single-chain surfactants the concentration of the single tail surfactants in all our simulations is well above its characteristic micellar concentration (CMC) while the concentration of the gemini surfactants is much lower. The total volume fraction ϕ is defined by $\phi = (N_{st}\ell + N_{gem}[2\ell + n])/(L_x L_y L_z)$ where N_{st} and N_{gem} are the total numbers of single-tail and gemini

surfactants, respectively, in the system. Moreover, the relative volume fractions of the geminis and the single-tail surfactants are defined by $c_{gem} = (N_{gem}[2\ell+n]) / (N_{st}\ell + N_{gem}[2\ell+n])$

We allow the same moves of the surfactants and water molecules as those allowed in ref. [18–20]. The moves allowed for the surfactants in our model are as follows:

(i) *reptation*: this is identical to the reptation move for single-chain surfactants (also called "slithering snake" move) [18–20]; (ii) *spontaneous chain buckling*: a portion in the middle of one of the two tails or the spacer is randomly picked up and allowed to buckle with the Boltzmann probability mentioned above; (iii) *kink movement*: a kink formed by the buckling or reptation is allowed to move to a new position with the appropriate Boltzmann probability; (iv) *pull move*: this is the reverse of spontaneous chain buckling; a buckled part of one of the two tails or the spacer is pulled so as to make it more extended. Each of these moves is possible only if the new positions of all the monomers are not occupied simultaneously by monomers belonging to other surfactants. Each surfactant is allowed to try each of the above mentioned moves once during each MC step. The system evolves following the standard Metropolis algorithm [25] while we monitor the mean-size of the clusters of the surfactants. After sufficiently long time, if the mean size of the clusters of the surfactants attains a time-independent value it indicates that the system has been equilibrated. All the results (except the snapshots) reported in this paper have been generated for system sizes $L_x = L_y = L_z = L = 100$ by averaging over sufficiently large (10-25) number of runs.

We have defined CMC as the amphiphile concentration where half of the surfactants are in the form of isolated chains and the other half in the form of clusters consisting of more than one neighbouring amphiphile [16,20].

To quantify the process of cross-linking we have calculated the number of cross-links in the system as follows: a) First we make a Hoshen-Kopelman [26] cluster count without the gemini surfactants i.e we identify and mark the cluster formed by the single tail surfactants. b) Next we do a second Hoshen-Kopelman cluster count with both gemini and single tail surfactants. c) When a cluster in (b) dissolves in two or more clusters in (a) we have found

a cross-linked micelle. The number of surfactant monomers per micelle is also counted by a Hoshen-Kopelman algorithm [26,27].

III. AGGREGATION NUMBERS

We have computed the aggregation numbers of micelles present in the gemini/single tail surfactants at different temperature and for several concentrations. Figure 3 and 4 show the variations of the aggregation number of the pure non-ionic gemini and single tail surfactants for different concentrations and temperatures. The aggregation number of the pure gemini micelles increases with concentration and temperature as for conventional non-ionic surfactants. For the single tail surfactants ($\mathcal{T}_{10}\mathcal{N}_1\mathcal{H}_1$) the mean aggregation number has the same behaviour as that of gemini surfactants. At a given temperature the aggregation number is higher for higher concentration of surfactants. This is in qualitative agreement with the previous experimental results [28]. Figure 5 shows the variation of N_A , the total number of surfactant monomers (gemini plus single tail) per micelle in the mixed system, at a fixed total surfactant concentration, as a function of the percentage of the gemini surfactant present in the mixture. It is evident from figure 5 that the addition of the gemini surfactants swells the micelle size.

IV. CROSS-LINKED MICELLES

Addition of the gemini surfactants have a profound effect on the mechanism of micellization and overall micellar morphology. A few of the different types of supra-molecular aggregates that are expected to be formed by a mixture of single-tail surfactants and geminis in water are shown schematically in fig.6. It is possible that the gemini bind in an intramicellar fashion shown in mixed micelle (figure 6(a)) and the spacer reside at the micelle surface. The micelles may also be cross-linked (figure 6(b)). Another possibility is that the gemini reside in the middle of the micellar aggregates.

An instantaneous snapshot of two spherical micelles cross-linked by a gemini surfactant is shown in fig.7; the two micelles consist, almost exclusively, of single-tail surfactants while the cross-linking is done by a gemini surfactant. However, an instantaneous snapshot of a more exotic supra-molecular aggregate is shown in fig.8; a "necklace"-like aggregate (with branch) is formed by the cross-linking of a sequence of micelles by gemini surfactants. These snapshots not only provide the most direct evidence in favour of cross-linking of micelles but also establish the formation of "aggregates of micellar aggregates" by the mixture of single-tail surfactants and geminis in water.

In our simulations of a mixture of "telechelic chain" and single-tail surfactants also, we get cross-linked micelles. However, the number of cross-links is less than that in the case when gemini surfactant is present. When a "telechelic chain" cross-links, the only gain in energy comes from the fact that their heads find themselves close to other heads of single-chain surfactants on the surface of the micelles. But, they do not have any tails to hide and therefore, not as much urge to cross-link as the geminis.

The cross-linking of the micelles is dynamic, rather than permanent, in the sense that such links can break while new links can form elsewhere in the system with time. To our knowledge this is the first direct evidence of the formation of cross-linked micelles. From our extensive simulations we find that they occur, for both flexible as well as rigid hydrophilic spacer, when the spacer is sufficiently long. When the spacer is hydrophilic there is the following scenario (figure 6):

- If $d_{mic} \gg$ typical extension of the spacer, case 6(a) is preferred. In that case we get swelling of micelles.
- If $d_{mic} \leq$ typical extension of the spacer, both cases 6(a) and 6(b) preferred.
- If $r_{mic} \ll$ typical extension of the spacer then cross-linking is preferred. For long spacer the correction in the entropic term is logarithmic, the cross-linking is weakly preferred.

Figure 9 shows the mean number of cross-linked micelles as a function of the fraction of

the gemini surfactants present in the mixture. At high gemini concentration geminis form their own micelles rather than cross-linking. By forming their own micelles geminis gain entropically. That is why the number of cross-linked micelles decreases when there is large percentage of geminis present in the system.

We have also computed the CMC for the mixtures of geminis and single tail surfactants. In Figure 10 we have plotted the CMC of the mixture against the fraction of gemini surfactants. The plot shows the presence of a minimum which indicates synergism in micelle formation in the mixture. Synergism in mixed micelle formation has been previously reported [29,30] for mixture of anionic gemini surfactants and various conventional single tail non-ionic surfactants.

We have also carried out simulations when the spacer is rigid. One of the very surprising observations is that when we have rigid spacer, we get crosslinking both for hydrophilic as well as hydrophobic spacers. However for hydrophobic spacers we observed fewer cross-linked micelles. Figure 11 shows a snapshot of the cross-linked micelle when the rigid spacer is hydrophobic. This is in qualitative agreement with the result of Menger et. al. [11] where they postulated the occurrence of micelle cross-linking in a mixture of CTAB and gemini surfactant with rigid spacer.

We have also carried out simulations when the length of the spacer $n = 0$; this corresponds to model lipids rather than geminis (see fig. 2). In this case, however, we do not find any cross-linked micelles. Instead, we observed *mixed micelles* which have been shown schematically in fig.6. Hydrophilic spacers gain energy by remaining surrounded by water. That is why in the snapshots of micellar aggregates we see that a larger number of monomers belonging to the spacers are in contact with water. We believe that the overall loss of entropy caused by the formation of "cross-linked" micelles cannot be compensated by the gain of energy if the hydrophilic spacers are not long enough. This explains why mixtures of single-tail surfactants and lipids were never seen to form cross-linked micelles in our simulations.

We have also studied quantitatively the growth of the micelles with time starting from

initial states where surfactants are dispersed randomly in water. Initially, we observe an increase of the mean-cluster size with the passage of time (see fig.12); this indicates the spontaneous formation and subsequent growth of the micellar aggregates. Interestingly, the mean-cluster size appears not only to follow a power-law growth but also displays three different regimes of growth, each characterized by a distinct growth exponent (fig.12) for extremely long times. For higher temperatures the growth stopped after some time.

The formation and growth of micelles from a random initial configuration take place during the first regime (0 – 14 million MCS). In the second regime (14 – 42 million MCS) the micelles start coalescing thereby forming an intermediate structure and this process is very fast. During the next regime (42 – 200 million MCS) this intermediate structure spontaneously forms vesicles which get cross-linked with either vesicles or micelles. Thus, from this unusually long runs we find a transition from micelles to vesicles. A snapshot showing the vesicles cross-linked with micelles is shown in figure 13. Figure 14 shows a cross-section of the vesicle. Our simulations might provide some insight into the kinetics and possible existence as well as structure of intermediates during the micelle-to-vesicles transition. Although the exponents are close to those associated with Ostwald ripening in binary alloys and spinodal decomposition in binary fluids [31], we leave it for future publications to explore this non-equilibrium growth process.

V. DISCUSSION

Here we try to make some connection between the available experimental parameters and our simulation parameters. For the snapshot shown in figure 7 the number of gemini surfactants is 32 and the number of single tail surfactant is 1200. Since the total lattice site is 10^6 , the mole fraction is 3.2×10^{-5} for geminis and 1.2×10^{-3} for single-tail surfactants. These mole fractions are in the same range of experimental mole fractions reported by Menger et. al [11] where they postulated the possibility of micelle cross-linking. In their experiments the mole fraction of the gemini was in the range 1.9×10^{-5} to 5.6×10^{-4} and

that for CTAB (single tail surfactants used in the experiment) was in the range 5.0×10^{-3} . Moreover, CMC of CTAB is 8×10^{-4} . The CMC of the single tail surfactants $\mathcal{T}_4\mathcal{N}_1\mathcal{H}_1$ used in our simulation is in the range 10^{-3} to 10^{-4} [17] which is comparable to the CMC of CTAB. These comparisons suggest that our choice of parameters is reasonable and our simulation results can be compared with the corresponding experimental results.

The present study gives direct evidence of the formation of cross-linked micelles in a mixture of single tail and gemini surfactants. When the spacer of the gemini is hydrophilic and sufficiently long cross-linking occurs. However when the spacer is hydrophobic, cross-linking occurs only when the spacer is completely rigid. Our simulation also elucidate the mechanism of mixed micellization. We have also calculated the aggregation numbers of the mixed micelles as a function of temperature as well as concentration. The variation of the CMC of the mixture shown synergism in micelle formation. From our longest simulation run we observe a micelle-vesicle transition. This might give us some understanding of the growth process and intermediate structure during micelle-vesicle transition. Our preliminary results in a mixture of telechelic chain and single tail surfactants also gives rise to cross-linking of micelles. Finally our simulations show how the micellar size and its morphology can be controlled by adding dimeric surfactants or telechelic chains.

Acknowledgements: We thank R. Zana for useful comments and B. Dünweg for helpful discussions. PKM and DC acknowledge financial supports from the Alexander von Humboldt foundation and SFB 341 Köln-Aachen-Jülich, respectively.

¹ Present address: Dept. of Physics, Condensed Matter Laboratory, University of Colorado, Boulder, CO 80309-0390, USA.

* Email : maiti@colorado.edu.

[†] On leave from Physics Dept., I.I.T., Kanpur, India.

- [1] Evans D. F. and Wennerstrom H., *The colloidal domain where physics, chemistry, biology and technology meet*, VCH, New York (1994).
- [2] Deinega Y. F., Ulberg Z. R., Marochko L. G., Rudi V. P. and Deisenko V. P., *Kolloidn Zh.* **36**, 649 (1974).
- [3] Menger F. M. and Littau C. A., *J. Am. Chem. Soc.*, **113**, 1451 (1991).
- [4] Menger F. M. and Littau C. A., *J. Am. Chem. Soc.*, **115**, 10083 (1993).
- [5] Rosen M., *Chemtech* **23**, 30 (1993).
- [6] Zana R., Benrraou M. and Rueff R., *Langmuir*, **7**, 1072 (1991).
- [7] Zana R. and Talmon Y., *Nature*, **362**, 228 (1993).
- [8] Zana R., in: *Novel Surfactants: Preparation, Applications and Biodegradability*, ed. Holmberg C (Dekker, New York, 1998)
- [9] Song L. D. and Rosen M. J., *Langmuir*, **12**, 1149 (1996).
- [10] Gelbart W., Ben Shaul A. and Roux D. (eds.) *Micelles, Membranes, Microemulsions and Monolayers* (Springer, Berlin, 1994). Tanford C., *The Hydrophobic Effect: Formation of Micelles and Biological Membranes*, (Wiley, New York 1980).
- [11] Menger F. M. and Eliseev A. V., *Langmuir*, **11**, 1855 (1995).
- [12] Zana R., Levy H., Danino D., Talmon Y. and Kwetkat K., *Langmuir*, **13**, 402 (1997).
- [13] Zana R., Levy H. and Kwetkat K., *J. Colloid Interface Sci.*, **197**, 370 (1998).
- [14] Khalatur, P. G. and Khokhlov, A. R., *Macromol. Theory Simul.*, **5**, 877, (1996).
- [15] Semenov, A. N., Joanny, J. F. and Khokhlov, A. R., *Macromolecules*, **28**, 1066 (1995).
- [16] Stauffer D., Jan N., He Y., Pandey R. B., Marangoni D. G. and Smith-Palmer T., *J. Chem. Phys.* **100**, 6934 (1994);

- [17] Jan N. and Stauffer D., *J. de Physique I* **4**, 345 (1994); D. Stauffer and D. Woermann, *J. Phys. II (France)*, **5**, 1 (1995).
- [18] Liverpool T. B., in: *Annual Reviews of Computational Physics*, vol. IV, ed. D. Stauffer (World Scientific, Singapore 1996). Schmid, F. in : *Computational methods in colloid and interface science*, ed. M. Borowko, (Marcel Dekker inc.), in press.
- [19] Bernardes A. T., *J. Phys. II France* **6**, 169 (1996); see also *Langmuir*, **12**, 5763 (1996).
- [20] Maiti P. K. and Chowdhury D., *Europhys. Lett.* **41**, 183, (1998); *J. Chem. Phys.* **109**, 5126 (1998)
- [21] Smit B., Hilbers P. A. J., Esselink K., Rupert L. A. M., van Os N. M. and Schlijper A. G., *Nature*, **348**, 624 (1990); Karaboni S., Esselink K., Hilbers P. A. J., Smit B., Karthaus J., van Os N. M. and Zana R., *Science*, **266**, 254 (1994).
- [22] Larson R. G., Scriven L. E. and Davis H. T., *J. Chem. Phys.* **83**, 2411 (1985); Larson R. G., *J. Chem. Phys.* **89**, 1642 (1988) and **91**, 2479 (1989).
- [23] Gompper G. and Schick M., *Phase Transitions and Critical Phenomena*, Vol. 16, ed. C. Domb and J. L. Lebowitz, (Academic Press, London 1994).
- [24] Chowdhury D., *Langmuir*, **12**, 1098 (1996); Chowdhury D., Maiti P. K., Sabhapandit S. and Taneja P., *Phys. Rev.E*, **56**, 667 (1997).
- [25] Binder K. (ed.) *Applications of the Monte Carlo Method in Statistical Physics*, Topics in Applied Physics 36, (Springer, 1987).
- [26] Hoshen J. and Kopelman R., *Phys. Rev.* **B14**, 3438 (1976); see also Stauffer D. and Aharony A., *Introduction to percolation theory*, (Taylor & Francis, London, 1992).
- [27] Flimm O., Masters Thesis, University of Cologne (unpublished, 1999)
- [28] Zana R. and Weill C., *J. Phys. Lett.* **46**, L953 (1985).

- [29] Rosen M. J., Gao T., Nakatsuji Y. and Masuyamai A., Colloids Surf. A: Physicochem. Eng. Asp. **88**, 1 (1994).
- [30] Rosen M. J., Zhu Z. and Gao T., J. Colloid Interface Sci., **15**, 224 (1993).
- [31] Gunton J. D. and Droz M., *Lec. Notes in Physics*, vol. 183 (1983) ; A.J. Bray, Adv. Phys.**43**, 357 (1994).






FIG. 1. A schematic representation of different types of amphiphiles.




Single Chain



Double Chain (Lipid)



Gemini with Spacer 2



Telechelic Chain

● Head ○ Tail/Spacer ⊕ Liaison

FIG. 2. Models of single-chain, double-chain, gemini surfactants and telechelic chain.




FIG. 3. Variation of micelle aggregation number N_A with temperature formed by gemini surfactants in water for gemini concentration $c_{gem} = 0.005(\circ)$; $0.01(\square)$; $0.02(\diamond)$; and $0.03(\triangle)$.




FIG. 4. Variation of micelle aggregation number N_A with temperature formed by single-tail surfactants in water for surfactant concentration $c_{st} = 0.005(\diamond)$; $0.013(+)$; and $0.022(\square)$. The continuous lines are merely guide to the eye.




FIG. 5. Variation of mixed micelle aggregation number N_A with fraction of gemini present in the mixture of single-tail and gemini surfactants in water for different temperature T and surfactant concentration ϕ . The symbols correspond to $T = 2.2(\square)$ and $3.0(\diamond)$ at $\phi = 0.022$ and $T = 2.2(\triangle)$ and $3.0(*)$ at $\phi = 0.013$.




FIG. 6. Schematic representation of a few possible supra-molecular aggregates formed by single-tail and gemini surfactants in water. r_{mic} is the size of the micelle and d_{mic} is the distance between the micelles.




FIG. 7. One of the cross-linked micelles in the system is shown. The parameters are: $m = 4$ for both single-tail surfactants and geminis and $n = 13$ for the geminis while $T = 2.2$, $\phi = 0.008$, $c_{gem} = 0.1$. The monomers belonging to the heads, neutral parts and tails of the single-tail surfactants are represented by red, gray and blue spheres while those belonging to the heads (as well as spacers), neutral parts and tails of the geminis are represented by yellow, white and green spheres, respectively.




FIG. 8. One necklace-like aggregate (with branch) formed by the cross-linking of micelles in the system is shown. The parameters are: $m = 8$ for the single-tail surfactants while $m = 4, n = 13$ for the geminis and $T = 2.8, \phi = 0.008, c_{gem} = 0.1$. The colour code is same as in figure 7.




FIG. 9. Percentage of cross-linked micelles as a function of the fraction of the gemini. The total concentration of the surfactants is 0.03 and temperature is 2.2.




FIG. 10. Variation of the cmc (total concentration of surfactant) of the gemini/single-tail surfactant mixture with gemini mole fraction at temperature $T = 2.2$.




FIG. 11. Cross-linked micelles for hydrophobic rigid spacer is shown. The parameters are: $m = 10$ for both single-tail surfactants and geminis and $n = 14$ for the geminis while $T = 2.2, \phi = 0.022, c_{gem} = 0.005$. The colour code is same as in figure 7.




FIG. 12. The mean-cluster size of the monomers of the amphiphiles is plotted against time.




FIG. 13. Vesicles cross-linked with micelles after 119 million MCS. The colour code is same as in figure 7.




FIG. 14. Cross-section of the vesicle. White sphere represents hydrophilic monomers and grey sphere represent hydrophobic monomers.